

## Product datasheet for RC221690L3V

**OriGene Technologies, Inc.**  
9620 Medical Center Drive, Ste 200  
Rockville, MD 20850, US  
Phone: +1-888-267-4436  
<https://www.origene.com>  
[techsupport@origene.com](mailto:techsupport@origene.com)  
EU: [info-de@origene.com](mailto:info-de@origene.com)  
CN: [techsupport@origene.cn](mailto:techsupport@origene.cn)

### DKK3 (NM\_001018057) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

**Product Type:** Lentiviral Particles

**Symbol:** DKK3

**Synonyms:** REIC; RIG

**Mammalian Cell:** Puromycin

**Selection:**

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

**Tag:** Myc-DDK

**ACCN:** NM\_001018057

**ORF Size:** 1050 bp

**ORF Nucleotide Sequence:** The ORF insert of this clone is exactly the same as (RC221690).

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**RefSeq:** [NM\\_001018057.1](#), [NP\\_001018067.1](#)

**RefSeq Size:** 2587 bp

**RefSeq ORF:** 1053 bp

**Locus ID:** 27122

**UniProt ID:** [Q9UBP4](#)

**Cytogenetics:** 11p15.3



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use.

©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

1 / 2

**Protein Families:** Druggable Genome, Secreted Protein

**MW:** 38.3 kDa

**Gene Summary:** This gene encodes a protein that is a member of the dickkopf family. The secreted protein contains two cysteine rich regions and is involved in embryonic development through its interactions with the Wnt signaling pathway. The expression of this gene is decreased in a variety of cancer cell lines and it may function as a tumor suppressor gene. Alternative splicing results in multiple transcript variants encoding the same protein. [provided by RefSeq, Jul 2008]