

Product datasheet for RC221717L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OXGR1 (NM_080818) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: OXGR1 (NM_080818) Human Tagged ORF Clone Lentiviral Particle

Symbol: OXGR1

Synonyms: aKGR; GPR80; GPR99; P2RY15; P2Y15

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM_080818

ORF Size: 1011 bp

ORF Nucleotide

OTI Disclaimer:

tide The

Sequence:

The ORF insert of this clone is exactly the same as(RC221717).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through paturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 080818.3

 RefSeq Size:
 2285 bp

 RefSeq ORF:
 1014 bp

 Locus ID:
 27199

 UniProt ID:
 Q96P68

 Cytogenetics:
 13q32.1

Domains: 7tm 1

Protein Families: Druggable Genome, GPCR, Transmembrane



ORIGENE

MW: 38.1 kDa

Gene Summary: This gene encodes a G protein-coupled receptor (GPCR) that belongs to the oxoglutarate

receptor family within the GPCR superfamily. The encoded protein is activated by the citric acid intermediate, oxoglutarate, as well as several cysteinyl leukotrienes, including leukotrienes E4, C4 and D4, which are implicated in many inflammatory disorders. In mice, a

knock-out of this gene leads to middle ear inflammation, changes in the mucosal epithelium, and an increase in fluid behind the eardrum, and is associated with hearing loss. Alternative

splicing results in multiple transcript variants. [provided by RefSeq, Oct 2016]