

Product datasheet for RC221737L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

beta Arrestin 1 (ARRB1) (NM 020251) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: beta Arrestin 1 (ARRB1) (NM_020251) Human Tagged ORF Clone Lentiviral Particle

Symbol: beta Arrestin 1

Synonyms: ARB1; ARR1

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_020251 **ORF Size:** 1230 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC221737).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 020251.2

RefSeq Size: 2180 bp
RefSeq ORF: 1233 bp
Locus ID: 408

 UniProt ID:
 P49407

 Cytogenetics:
 11q13.4

Protein Families: Druggable Genome

Protein Pathways: Chemokine signaling pathway, Endocytosis, MAPK signaling pathway





beta Arrestin 1 (ARRB1) (NM_020251) Human Tagged ORF Clone Lentiviral Particle – RC221737L4V

MW: 46.1 kDa

Gene Summary:

Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of G-protein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 1 is a cytosolic protein and acts as a cofactor in the beta-adrenergic receptor kinase (BARK) mediated desensitization of beta-adrenergic receptors. Besides the central nervous system, it is expressed at high levels in peripheral blood leukocytes, and thus the BARK/beta-arrestin system is believed to play a major role in regulating receptor-mediated immune functions. Alternatively spliced transcripts encoding different isoforms of arrestin beta 1 have been described. [provided by RefSeq, Jan 2011]