

## Product datasheet for **RC221766L2V**

### EIF3A (NM\_003750) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	EIF3A (NM_003750) Human Tagged ORF Clone Lentiviral Particle
Symbol:	EIF3A
Synonyms:	EIF3; eIF3-p170; eIF3-theta; EIF3S10; P167; p180; p185; TIF32
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_003750
ORF Size:	4146 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC221766).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_003750.1</a> , <a href="#">NP_003741.1</a>
RefSeq Size:	5256 bp
RefSeq ORF:	4149 bp
Locus ID:	8661
UniProt ID:	<a href="#">Q14152</a>
Cytogenetics:	10q26.11
Domains:	PCI
MW:	166.4 kDa



[View online »](#)

**Gene Summary:**

RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:17581632, PubMed:25849773). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA<sub>i</sub> and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:17581632, PubMed:11169732). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773, PubMed:27462815). [UniProtKB/Swiss-Prot Function]