

Product datasheet for RC221789L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL28 Receptor alpha (IFNLR1) (NM 173065) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IL28 Receptor alpha (IFNLR1) (NM_173065) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL28 Receptor alpha

Synonyms: CRF2/12; IFNLR; IL-28R1; IL28RA; LICR2

Mammalian Cell

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 173065

Tag: Myc-DDK

ORF Size: 732 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC221789).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 173065.2

 RefSeq Size:
 4432 bp

 RefSeq ORF:
 735 bp

 Locus ID:
 163702

 UniProt ID:
 Q8IU57

 Cytogenetics:
 1p36.11

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway





IL28 Receptor alpha (IFNLR1) (NM_173065) Human Tagged ORF Clone Lentiviral Particle – RC221789L3V

MW: 27.5 kDa

Gene Summary:

The protein encoded by this gene belongs to the class II cytokine receptor family. This protein forms a receptor complex with interleukine 10 receptor, beta (IL10RB). The receptor complex has been shown to interact with three closely related cytokines, including interleukin 28A (IL28A), interleukin 28B (IL28B), and interleukin 29 (IL29). The expression of all three cytokines can be induced by viral infection. The cells overexpressing this protein have been found to have enhanced responses to IL28A and IL29, but decreased response to IL28B. Three alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]