

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC221836L4V

Optineurin (OPTN) (NM_001008213) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Optineurin (OPTN) (NM_001008213) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Optineurin
Synonyms:	ALS12; FIP2; GLC1E; HIP7; HYPL; NRP; TFIIIA-INTP
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001008213
ORF Size:	1731 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC221836).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001008213.1, NP 001008214.1</u>
RefSeq Size:	3582 bp
RefSeq ORF:	1734 bp
Locus ID:	10133
UniProt ID:	<u>Q96CV9</u>
Cytogenetics:	10p13
Protein Families:	Druggable Genome
MW:	65.9 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Optineurin (OPTN) (NM_001008213) Human Tagged ORF Clone Lentiviral Particle – RC221836L4V
Gene Summary:	This gene encodes the coiled-coil containing protein optineurin. Optineurin may play a role in normal-tension glaucoma and adult-onset primary open angle glaucoma. Optineurin
	interacts with adenovirus E3-14.7K protein and may utilize tumor necrosis factor-alpha or
	Fas-ligand pathways to mediate apoptosis, inflammation or vasoconstriction. Optineurin may

also function in cellular morphogenesis and membrane trafficking, vesicle trafficking, and transcription activation through its interactions with the RAB8, huntingtin, and transcription factor IIIA proteins. Alternative splicing results in multiple transcript variants encoding the same protein. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US