

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC221895L4V

PCB (PC) (NM_000920) Human Tagged ORF Clone Lentiviral Particle

Product data:

| Product Type: | Lentiviral Particles |
|------------------------------|---|
| Product Name: | PCB (PC) (NM_000920) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | PCB |
| Synonyms: | PCB |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-mGFP-P2A-Puro (PS100093) |
| Tag: | mGFP |
| ACCN: | NM_000920 |
| ORF Size: | 3534 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC221895). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | <u>NM 000920.2</u> |
| RefSeq Size: | 4111 bp |
| RefSeq ORF: | 3537 bp |
| Locus ID: | 5091 |
| UniProt ID: | <u>P11498</u> |
| Cytogenetics: | 11q13.2 |
| Protein Families: | Druggable Genome |
| Protein Pathways: | Citrate cycle (TCA cycle), Metabolic pathways, Pyruvate metabolism |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| | PCB (PC) (NM_000920) Human Tagged ORF Clone Lentiviral Particle – RC221895L4V |
|---------------|--|
| MW: | 129.63 kDa |
| Gene Summary: | This gene encodes pyruvate carboxylase, which requires biotin and ATP to catalyse the carboxylation of pyruvate to oxaloacetate. The active enzyme is a homotetramer arranged in a tetrahedron which is located exclusively in the mitochondrial matrix. Pyruvate carboxylase is involved in gluconeogenesis, lipogenesis, insulin secretion and synthesis of the neurotransmitter glutamate. Mutations in this gene have been associated with pyruvate carboxylate carboxylase deficiency. Alternatively spliced transcript variants with different 5' UTRs, but encoding the same protein, have been found for this gene. [provided by RefSeq, Jul 2008] |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US