

Product datasheet for RC221986L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TLR8 (NM_138636) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TLR8 (NM_138636) Human Tagged ORF Clone Lentiviral Particle

Symbol: TLR8
Synonyms: CD288
Mammalian Cell None

Selection:

CII

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_138636 **ORF Size:** 3123 bp

ORF Nucleotide

•

Sequence:

The ORF insert of this clone is exactly the same as(RC221986).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 138636.2

 RefSeq Size:
 4211 bp

 RefSeq ORF:
 3126 bp

 Locus ID:
 51311

 UniProt ID:
 Q9NR97

 Cytogenetics:
 Xp22.2

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Toll-like receptor signaling pathway





ORIGENE

MW: 119.6 kDa

Gene Summary:

The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This gene is predominantly expressed in lung and peripheral blood leukocytes, and lies in close proximity to another family member, TLR7, on chromosome X. [provided by RefSeq, Jul 2008]