

## Product datasheet for RC222039L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CDC2L1 (CDK11B) (NM 033487) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: CDC2L1 (CDK11B) (NM\_033487) Human Tagged ORF Clone Lentiviral Particle

Symbol: CDK11B

**Synonyms:** CDC2L1; CDK11-p46; CDK11-p58; CDK11-p110; CLK-1; p58; p58CDC2L1; p58CLK-1;

PITSLREA; PK58

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_033487 **ORF Size:** 1578 bp

**ORF Nucleotide** 

•

Sequence:

The ORF insert of this clone is exactly the same as(RC222039).

**OTI Disclaimer:** 

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 033487.1</u>

 RefSeq Size:
 2226 bp

 RefSeq ORF:
 1581 bp

 Locus ID:
 984

 UniProt ID:
 P21127

 Cytogenetics:
 1p36.33

**Domains:** pkinase, TyrKc, S\_TKc





## CDC2L1 (CDK11B) (NM\_033487) Human Tagged ORF Clone Lentiviral Particle - RC222039L4V

**Protein Families:** Druggable Genome, Transcription Factors

MW: 59.1 kDa

Gene Summary: This gene encodes a member of the serine/threonine protein kinase family. Members of this

kinase family are known to be essential for eukaryotic cell cycle control. Due to a segmental duplication, this gene shares very high sequence identity with a neighboring gene. These two genes are frequently deleted or altered in neuroblastoma. The protein kinase encoded by this gene can be cleaved by caspases and may play a role in cell apoptosis. Alternative

splicing results in multiple transcript variants. [provided by RefSeq, Apr 2014]