

## Product datasheet for RC222068L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## DNMT3B (NM\_175850) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: DNMT3B (NM 175850) Human Tagged ORF Clone Lentiviral Particle

Symbol: DNMT3E

Synonyms: ICF; ICF1; M.HsallIB

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 175850

ORF Size: 2535 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC222068).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 175850.2, NP 787046.1

RefSeq Size: 4272 bp
RefSeq ORF: 2538 bp
Locus ID: 1789
UniProt ID: Q9UBC3

Cytogenetics: 20q11.21

Protein Families: Druggable Genome, Embryonic stem cells, Induced pluripotent stem cells, Stem cell -

Pluripotency





## DNMT3B (NM\_175850) Human Tagged ORF Clone Lentiviral Particle - RC222068L3V

**Protein Pathways:** Cysteine and methionine metabolism, Metabolic pathways

**MW:** 94.7 kDa

Gene Summary: CpG methylation is an epigenetic modification that is important for embryonic development,

imprinting, and X-chromosome inactivation. Studies in mice have demonstrated that DNA

methylation is required for mammalian development. This gene encodes a DNA methyltransferase which is thought to function in de novo methylation, rather than

maintenance methylation. The protein localizes primarily to the nucleus and its expression is developmentally regulated. Mutations in this gene cause the immunodeficiency-centromeric instability-facial anomalies (ICF) syndrome. Eight alternatively spliced transcript variants have been described. The full length sequences of variants 4 and 5 have not been determined.

[provided by RefSeq, May 2011]