

# **Product datasheet for RC222078**

## VEGFA (NM 001025367) Human Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** VEGFA (NM\_001025367) Human Tagged ORF Clone

Tag: Myc-DDK Symbol: VEGFA

Synonyms: MVCD1; VEGF; VPF

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

ORF Nucleotide >RC222078 representing NM\_001025367
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

TGGACGCGGCGAGCCGCGGGCAGGGCCGGAGCCCGCGCGGAGGCGGGTGGAGGGGGTCGGGGC TCGCGGCGTCGCACTGAAACTTTTCGTCCAACTTCTGGGCTGTTCTCGCTTCGGAGGAGCCGTGGTCCGC GCGGGGGAAGCCGAGCCGAGCCGCAGCCGCAGAAGTGCTAGCTCGGGCCGGGAGGAGCCGCAGCCGGAGG AGGGGGAGGAGGAGAAGAAGAAGAAGAGAGGGGGGCCGCAGTGGCGACTCGGCGCTCGGAAGCCGGG CGGGCCTCGGGCCGGGGAGGAAGAGTAGCTCGCCGAGGCGCCCGAGGAGAGCGGCCCCCCACAGCCCGA GCCGGAGAGGGAGCCGCGCCCGGCCCGGTCGGGCCTCCGAAACCATGAACTTTCTGCTGTCTTG GGTGCATTGGAGCCTTGCCTGCTGCTCCTACCTCCACCATGCCAAGTGGTCCCAGGCTGCACCCATGGCA GAAGGAGGAGGCAGAATCATCACGAAGTGGTGAAGTTCATGGATGTCTATCAGCGCAGCTACTGCCATC CAATCGAGACCCTGGTGGACATCTTCCAGGAGTACCCTGATGAGATCGAGTACATCTTCAAGCCATCCTG TGTGCCCCTGATGCGATGCGGGGGCTGCTGCAATGACGAGGGCCTGGAGTGTGTGCCCACTGAGGAGTCC AACATCACCATGCAGATTATGCGGATCAAACCTCACCAAGGCCAGCACATAGGAGAGATGAGCTTCCTAC GGGAAAGGGGCAAAAACGAAAGCGCAAGAAATCCCGTCCCTGTGGGCCTTGCTCAGAGCGGAGAAAGCAT TTGTTTGTACAAGATCCGCAGACGTGTAAATGTTCCTGCAAAAACACAGACTCGCGTTGCAAGGCGAGGC AGCTTGAGTTAAACGAACGTACTTGCAGATGTGACAAGCCGAGGCGG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >RC222078 representing NM\_001025367

Red=Cloning site Green=Tags(s)

LTDRQTDTAPSPSYHLLPGRRRTVDAAASRGQGPEPAPGGGVEGVGARGVALKLFVQLLGCSRFGGAVVR AGEAEPSGAARSASSGREEPQPEEGEEEEKEEERGPQWRLGARKPGSWTGEAAVCADSAPAARAPQALA RASGRGGRVARRGAEESGPPHSPSRRGSASRAGPGRASETMNFLLSWVHWSLALLLYLHHAKWSQAAPMA EGGGQNHHEVVKFMDVYQRSYCHPIETLVDIFQEYPDEIEYIFKPSCVPLMRCGGCCNDEGLECVPTEES NITMQIMRIKPHQGQHIGEMSFLQHNKCECRPKKDRARQEKKSVRGKGKGQKRKRKKSRPCGPCSERRKH LFVQDPQTCKCSCKNTDSRCKARQLELNERTCRCDKPRR


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** 

Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the OR

**ACCN:** NM\_001025367

ORF Size: 1167 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

**RefSeq:** <u>NM 001025367.3</u>

 RefSeq Size:
 3608 bp

 RefSeq ORF:
 1170 bp

 Locus ID:
 7422

 UniProt ID:
 P15692

 Cytogenetics:
 6p21.1

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Bladder cancer, Cytokine-cytokine receptor interaction, Focal adhesion, mTOR signaling

pathway, Pancreatic cancer, Pathways in cancer, Renal cell carcinoma, VEGF signaling pathway


MW: 42.8 kDa



#### **Gene Summary:**

This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative inframe translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site. The levels of VEGF are increased during infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), thus promoting inflammation by facilitating recruitment of inflammatory cells, and by increasing the level of angiopoietin II (Ang II), one of two products of the SARS-CoV-2 binding target, angiotensin-converting enzyme 2 (ACE2). In turn, Ang II facilitates the elevation of VEGF, thus forming a vicious cycle in the release of inflammatory cytokines. [provided by RefSeq, Jun 2020]

### **Product images:**



Circular map for RC222078