

## Product datasheet for RC222128L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TECK (CCL25) (NM\_005624) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: TECK (CCL25) (NM\_005624) Human Tagged ORF Clone Lentiviral Particle

Symbol: TECK

**Synonyms:** Ckb15; Ck beta-15; SCYA25; TECK

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 005624

ORF Size: 450 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC222128).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 005624.2, NP 005615.2

 RefSeq Size:
 1002 bp

 RefSeq ORF:
 453 bp

 Locus ID:
 6370

 UniProt ID:
 015444

 Cytogenetics:
 19p13.2

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction





ORÏGENE

**MW:** 16.6 kDa

**Gene Summary:** This antimicrobial gene belongs to the subfamily of small cytokine CC genes. Cytokines are a

family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by

this gene displays chemotactic activity for dendritic cells, thymocytes, and activated

macrophages but is inactive on peripheral blood lymphocytes and neutrophils. The product of this gene binds to chemokine receptor CCR9. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Sep 2014]