

## Product datasheet for RC222275L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL28 Receptor alpha (IFNLR1) (NM 170743) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** IL28 Receptor alpha (IFNLR1) (NM\_170743) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL28 Receptor alpha

CRF2/12; IFNLR; IL-28R1; IL28RA; LICR2 Synonyms:

**Mammalian Cell** 

Selection:

ACCN:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK NM 170743

**ORF Size:** 1560 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC222275).

Sequence: OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 170743.2

RefSeq Size: 4549 bp RefSeq ORF: 1563 bp Locus ID: 163702 **UniProt ID:** Q8IU57 Cytogenetics: 1p36.11

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway





IL28 Receptor alpha (IFNLR1) (NM\_170743) Human Tagged ORF Clone Lentiviral Particle – RC222275L1V

**MW:** 57.5 kDa

**Gene Summary:** 

The protein encoded by this gene belongs to the class II cytokine receptor family. This protein forms a receptor complex with interleukine 10 receptor, beta (IL10RB). The receptor complex has been shown to interact with three closely related cytokines, including interleukin 28A (IL28A), interleukin 28B (IL28B), and interleukin 29 (IL29). The expression of all three cytokines can be induced by viral infection. The cells overexpressing this protein have been found to have enhanced responses to IL28A and IL29, but decreased response to IL28B. Three alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]