

Product datasheet for RC222354L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PDX1 (NM_000209) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PDX1 (NM_000209) Human Tagged ORF Clone Lentiviral Particle

Symbol: PDX²

Synonyms: GSF; IDX-1; IPF1; IUF1; MODY4; PAGEN1; PDX-1; STF-1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 000209

ORF Size: 849 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC222354).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000209.1

 RefSeq Size:
 1525 bp

 RefSeq ORF:
 852 bp

 Locus ID:
 3651

 UniProt ID:
 P52945

 Cytogenetics:
 13q12.2

Protein Families: Embryonic stem cells, ES Cell Differentiation/IPS, Induced pluripotent stem cells, Transcription

Factors





PDX1 (NM_000209) Human Tagged ORF Clone Lentiviral Particle - RC222354L3V

Protein Pathways: Maturity onset diabetes of the young, Type II diabetes mellitus

MW: 30.6 kDa

Gene Summary: The protein encoded by this gene is a transcriptional activator of several genes, including

insulin, somatostatin, glucokinase, islet amyloid polypeptide, and glucose transporter type 2. The encoded nuclear protein is involved in the early development of the pancreas and plays a major role in glucose-dependent regulation of insulin gene expression. Defects in this gene are a cause of pancreatic agenesis, which can lead to early-onset insulin-dependent diabetes mellitus (IDDM), as well as maturity onset diabetes of the young type 4 (MODY4). [provided

by RefSeq, Aug 2017]