

## Product datasheet for RC222382L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# PDI (PDIA2) (NM\_006849) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** PDI (PDIA2) (NM\_006849) Human Tagged ORF Clone Lentiviral Particle

Symbol: PDI

Synonyms: PDA2; PDI; PDIP; PDIR

**Mammalian Cell** 

Selection:

ACCN:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 006849

Tag: Myc-DDK

ORF Size: 1575 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC222382).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 006849.2</u>

 RefSeq Size:
 1726 bp

 RefSeq ORF:
 1578 bp

 Locus ID:
 64714

 UniProt ID:
 Q13087

 Cytogenetics:
 16p13.3

**Protein Families:** Druggable Genome

**MW:** 58 kDa







#### **Gene Summary:**

This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, two catalytically active thioredoxin (TRX) domains, two TRX-like domains and a C-terminal ER-retention sequence. The protein plays a role in the folding of nascent proteins in the endoplasmic reticulum by forming disulfide bonds through its thiol isomerase, oxidase, and reductase activity. The encoded protein also possesses estradiol-binding activity and can modulate intracellular estradiol levels. [provided by RefSeq, Sep 2017]