

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC222382L4V

PDI (PDIA2) (NM_006849) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	PDI (PDIA2) (NM_006849) Human Tagged ORF Clone Lentiviral Particle
Symbol:	PDI
Synonyms:	PDA2; PDI; PDIP; PDIR
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_006849
ORF Size:	1575 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC222382).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 006849.2</u>
RefSeq Size:	1726 bp
RefSeq ORF:	1578 bp
Locus ID:	64714
UniProt ID:	<u>Q13087</u>
Cytogenetics:	16p13.3
Protein Families:	Druggable Genome
MW:	58 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic
reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions.
The encoded protein has an N-terminal ER-signal sequence, two catalytically active
thioredoxin (TRX) domains, two TRX-like domains and a C-terminal ER-retention sequence.
The protein plays a role in the folding of nascent proteins in the endoplasmic reticulum by
forming disulfide bonds through its thiol isomerase, oxidase, and reductase activity. The
encoded protein also possesses estradiol-binding activity and can modulate intracellular
estradiol levels. [provided by RefSeq, Sep 2017]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US