

## Product datasheet for RC222395L1V

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# DGKA (NM\_001345) Human Tagged ORF Clone Lentiviral Particle

### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** DGKA (NM\_001345) Human Tagged ORF Clone Lentiviral Particle

Symbol: DGKA

Synonyms: DAGK; DAGK1; DGK-alpha

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 001345

ORF Size: 2205 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC222395).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001345.4</u>

 RefSeq Size:
 2756 bp

 RefSeq ORF:
 2208 bp

 Locus ID:
 1606

 UniProt ID:
 P23743

 Cytogenetics:
 12q13.2

**Domains:** DAGKa, DAGKc, EFh, DAG\_PE-bind

**Protein Families:** Druggable Genome





### DGKA (NM\_001345) Human Tagged ORF Clone Lentiviral Particle - RC222395L1V

**Protein Pathways:** Glycerolipid metabolism, Glycerophospholipid metabolism, Metabolic pathways,

Phosphatidylinositol signaling system

**MW:** 82.5 kDa

**Gene Summary:** The protein encoded by this gene belongs to the eukaryotic diacylglycerol kinase family. It

acts as a modulator that competes with protein kinase C for the second messenger diacylglycerol in intracellular signaling pathways. It also plays an important role in the resynthesis of phosphatidylinositols and phosphorylating diacylglycerol to phosphatidic acid. Several transcript variants encoding different isoforms have been identified for this gene.

[provided by RefSeq, Apr 2017]