

## Product datasheet for RC222432L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PDGF beta (PDGFB) (NM\_033016) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** PDGF beta (PDGFB) (NM\_033016) Human Tagged ORF Clone Lentiviral Particle

Symbol: PDGF beta

**Synonyms:** c-sis; IBGC5; PDGF-2; PDGF2; SIS; SSV

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_033016

ORF Size: 678 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC222432).

Sequence:

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 033016.1, NP 148937.1

 RefSeq Size:
 2378 bp

 RefSeq ORF:
 681 bp

 Locus ID:
 5155

 UniProt ID:
 P01127

Cytogenetics: 22q13.1

**Protein Families:** Druggable Genome





## PDGF beta (PDGFB) (NM\_033016) Human Tagged ORF Clone Lentiviral Particle - RC222432L4V

**Protein Pathways:** Cytokine-cytokine receptor interaction, Focal adhesion, Gap junction, Glioma, MAPK signaling

pathway, Melanoma, Pathways in cancer, Prostate cancer, Regulation of actin cytoskeleton,

Renal cell carcinoma

MW: 25.3 kDa

**Gene Summary:** This gene encodes a member of the protein family comprised of both platelet-derived growth

factors (PDGF) and vascular endothelial growth factors (VEGF). The encoded preproprotein is proteolytically processed to generate platelet-derived growth factor subunit B, which can homodimerize, or alternatively, heterodimerize with the related platelet-derived growth factor subunit A. These proteins bind and activate PDGF receptor tyrosine kinases, which play a role in a wide range of developmental processes. Mutations in this gene are associated with meningioma. Reciprocal translocations between chromosomes 22 and 17, at sites where this

gene and that for collagen type 1, alpha 1 are located, are associated with

dermatofibrosarcoma protuberans, a rare skin tumor. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Oct 2015]