

Product datasheet for RC222603L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LRRC8C (NM_032270) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: LRRC8C (NM_032270) Human Tagged ORF Clone Lentiviral Particle

Symbol: LRRC80

Synonyms: AD158; FAD158

Mammalian Cell

Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 032270

ORF Size: 2409 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC222603).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 032270.2

 RefSeq Size:
 7207 bp

 RefSeq ORF:
 2412 bp

 Locus ID:
 84230

 UniProt ID:
 Q8TDW0

Cytogenetics: 1p22.2

Domains: LRR, LRR_TYP, LRR_PS

Protein Families: Transmembrane





ORIGENE

MW: 92.4 kDa

Gene Summary:

Non-essential component of the volume-regulated anion channel (VRAC, also named VSOAC channel), an anion channel required to maintain a constant cell volume in response to extracellular or intracellular osmotic changes. The VRAC channel conducts iodide better than chloride and can also conduct organic osmolytes like taurine. Plays a redundant role in the efflux of amino acids, such as aspartate and glutamate, in response to osmotic stress. Channel activity requires LRRC8A plus at least one other family member (LRRC8B, LRRC8C, LRRC8D or LRRC8E); channel characteristics depend on the precise subunit composition. [UniProtKB/Swiss-Prot Function]