

Product datasheet for RC222681L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GALNT11 (NM_022087) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: GALNT11 (NM_022087) Human Tagged ORF Clone Lentiviral Particle

Symbol: GALNT11

Synonyms: GALNAC-T11; GALNACT11

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_022087 **ORF Size:** 1824 bp

ORF Nucleotide

OTI Disclaimer:

102100

Sequence:

The ORF insert of this clone is exactly the same as(RC222681).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 022087.2

 RefSeq Size:
 2732 bp

 RefSeq ORF:
 1827 bp

 Locus ID:
 63917

 UniProt ID:
 Q8NCW6

 Cytogenetics:
 7q36.1

Domains: RICIN, Glycos_transf_2

Protein Families: Transmembrane





GALNT11 (NM_022087) Human Tagged ORF Clone Lentiviral Particle - RC222681L4V

Protein Pathways: Metabolic pathways, O-Glycan biosynthesis

MW: 68.9 kDa

Gene Summary: Polypeptide N-acetylgalactosaminyltransferase that catalyzes the initiation of protein O-

linked glycosylation and is involved in left/right asymmetry by mediating O-glycosylation of NOTCH1. O-glycosylation of NOTCH1 promotes activation of NOTCH1, modulating the balance between motile and immotile (sensory) cilia at the left-right organiser (LRO). Polypeptide N-acetylgalactosaminyltransferases catalyze the transfer of an N-acetyl-D-galactosamine residue to a serine or threonine residue on the protein receptor. Displays the

same enzyme activity toward MUC1, MUC4, and EA2 than GALNT1. Not involved in

glycosylation of erythropoietin (EPO).[UniProtKB/Swiss-Prot Function]