

Product datasheet for RC222696L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Glucose Transporter GLUT1 (SLC2A1) (NM_006516) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Glucose Transporter GLUT1 (SLC2A1) (NM_006516) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Glucose Transporter GLUT1

Synonyms: CSE; DYT9; DYT17; DYT18; EIG12; GLUT; GLUT-1; GLUT1DS; HTLVR; PED; SDCHCN

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_006516 **ORF Size:** 1476 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC222696).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 006516.1</u>

 RefSeq Size:
 2856 bp

 RefSeq ORF:
 1479 bp

 Locus ID:
 6513

 UniProt ID:
 P11166

 Cytogenetics:
 1p34.2

Domains: sugar_tr





Glucose Transporter GLUT1 (SLC2A1) (NM_006516) Human Tagged ORF Clone Lentiviral Particle – RC222696L4V

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Adipocytokine signaling pathway, Pathways in cancer, Renal cell carcinoma

MW: 53.9 kDa

Gene Summary: This gene encodes a major glucose transporter in the mammalian blood-brain barrier. The

encoded protein is found primarily in the cell membrane and on the cell surface, where it can also function as a receptor for human T-cell leukemia virus (HTLV) I and II. Mutations in this gene have been found in a family with paroxysmal exertion-induced dyskinesia. [provided by

RefSeq, Apr 2013]