

Product datasheet for **RC222709L4V**

ATF 4 (ATF4) (NM_001675) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	ATF 4 (ATF4) (NM_001675) Human Tagged ORF Clone Lentiviral Particle
Symbol:	ATF4
Synonyms:	CREB-2; CREB2; TAXREB67; TXREB
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001675
ORF Size:	1053 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC222709).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001675.2
RefSeq Size:	2022 bp
RefSeq ORF:	1056 bp
Locus ID:	468
UniProt ID:	P18848
Cytogenetics:	22q13.1
Domains:	BRLZ
Protein Families:	Transcription Factors



[View online »](#)

Protein Pathways:	GnRH signaling pathway, Long-term potentiation, MAPK signaling pathway, Neurotrophin signaling pathway, Prostate cancer
MW:	38.4 kDa
Gene Summary:	<p>This gene encodes a transcription factor that was originally identified as a widely expressed mammalian DNA binding protein that could bind a tax-responsive enhancer element in the LTR of HTLV-1. The encoded protein was also isolated and characterized as the cAMP-response element binding protein 2 (CREB-2). The protein encoded by this gene belongs to a family of DNA-binding proteins that includes the AP-1 family of transcription factors, cAMP-response element binding proteins (CREBs) and CREB-like proteins. These transcription factors share a leucine zipper region that is involved in protein-protein interactions, located C-terminal to a stretch of basic amino acids that functions as a DNA binding domain. Two alternative transcripts encoding the same protein have been described. Two pseudogenes are located on the X chromosome at q28 in a region containing a large inverted duplication. [provided by RefSeq, Sep 2011]</p>