

## **Product datasheet for RC222719**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## Cathepsin B (CTSB) (NM\_147780) Human Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: Cathepsin B (CTSB) (NM\_147780) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: Cathepsin B

Synonyms: APPS; CPSB; RECEUP

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >RC222719 representing NM\_147780

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA





**Protein Sequence:** >RC222719 representing NM\_147780

Red=Cloning site Green=Tags(s)

MWQLWASLCCLLVLANARSRPSFHPLSDELVNYVNKRNTTWQAGHNFYNVDMSYLKRLCGTFLGGPKPPQ RVMFTEDLKLPASFDAREQWPQCPTIKEIRDQGSCGSCWAFGAVEAISDRICIHTNAHVSVEVSAEDLLT CCGSMCGDGCNGGYPAEAWNFWTRKGLVSGGLYESHVGCRPYSIPPCEHHVNGSRPPCTGEGDTPKCSKI CEPGYSPTYKQDKHYGYNSYSVSNSEKDIMAEIYKNGPVEGAFSVYSDFLLYKSGVYQHVTGEMMGGHAI RILGWGVENGTPYWLVANSWNTDWGDNGFFKILRGQDHCGIESEVVAGIPRTDQYWEKI


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk8076">https://cdn.origene.com/chromatograms/mk8076</a> c03.zip

**Restriction Sites:** Sgfl-Mlul

Cloning Scheme:





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_147780

ORF Size: 1017 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 147780.4</u>

 RefSeq Size:
 3945 bp

 RefSeq ORF:
 1020 bp

 Locus ID:
 1508

 UniProt ID:
 P07858

 Cytogenetics:
 8p23.1

**Protein Families:** Druggable Genome, Protease

**Protein Pathways:** Antigen processing and presentation, Lysosome

**MW:** 35.9 kDa

**Gene Summary:** This gene encodes a member of the C1 family of peptidases. Alternative splicing of this gene

results in multiple transcript variants. At least one of these variants encodes a preproprotein that is proteolytically processed to generate multiple protein products. These products include the cathepsin B light and heavy chains, which can dimerize to form the double chain form of the enzyme. This enzyme is a lysosomal cysteine protease with both endopeptidase and exopeptidase activity that may play a role in protein turnover. It is also known as amyloid


precursor protein secretase and is involved in the proteolytic processing of amyloid

precursor protein (APP). Incomplete proteolytic processing of APP has been suggested to be a causative factor in Alzheimer's disease, the most common cause of dementia. Overexpression of the encoded protein has been associated with esophageal adenocarcinoma and other tumors. Both Cathepsin B and Cathepsin L are involved in the cleavage of the spike protein from the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) upon its entry to the human host cell. Multiple pseudogenes of this gene have been identified. [provided by


RefSeq, Sep 2020]



## **Product images:**



Circular map for RC222719



Western blot validation of overexpression lysate (Cat# [LY407800]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC222719 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).