

Product datasheet for **RC222757L1V**

Kv4.3 (KCND3) (NM_004980) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Kv4.3 (KCND3) (NM_004980) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Kv4.3
Synonyms:	BRGDA9; KCND3L; KCND3S; KSHIVB; KV4.3; SCA19; SCA22
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_004980
ORF Size:	1965 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC222757).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_004980.3
RefSeq Size:	2693 bp
RefSeq ORF:	1968 bp
Locus ID:	3752
UniProt ID:	Q9UK17
Cytogenetics:	1p13.2
Protein Families:	Druggable Genome, Ion Channels: Potassium, Transmembrane
MW:	73.3 kDa



[View online »](#)

Gene Summary:

Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. Four sequence-related potassium channel genes - shaker, shaw, shab, and shal - have been identified in *Drosophila*, and each has been shown to have human homolog(s). This gene encodes a member of the potassium channel, voltage-gated, shal-related subfamily, members of which form voltage-activated A-type potassium ion channels and are prominent in the repolarization phase of the action potential. This member includes two isoforms with different sizes, which are encoded by alternatively spliced transcript variants of this gene. [provided by RefSeq, Jul 2008]