

Product datasheet for RC222845L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Antithrombin III (SERPINC1) (NM 000488) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Antithrombin III (SERPINC1) (NM_000488) Human Tagged ORF Clone Lentiviral Particle

Symbol: Antithrombin III

Synonyms: AT3; AT3D; ATIII; ATIII-R2; ATIII-T1; ATIII-T2; THPH7

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 000488

ORF Size: 1392 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC222845).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000488.2

RefSeq Size: 1559 bp
RefSeq ORF: 1395 bp
Locus ID: 462

Locus ID: 462 **UniProt ID:** P01008

Cytogenetics: 1q25.1

Domains: SERPIN

Protein Families: Druggable Genome, Secreted Protein





Antithrombin III (SERPINC1) (NM_000488) Human Tagged ORF Clone Lentiviral Particle – RC222845L3V

Protein Pathways: Complement and coagulation cascades

MW: 52.6 kDa

Gene Summary: The protein encoded by this gene, antithrombin III, is a plasma protease inhibitor and a

member of the serpin superfamily. This protein inhibits thrombin as well as other activated serine proteases of the coagulation system, and it regulates the blood coagulation cascade. The protein includes two functional domains: the heparin binding-domain at the N-terminus of the mature protein, and the reactive site domain at the C-terminus. The inhibitory activity is enhanced by the presence of heparin. Numerous mutations have been identified for this gene, many of which are known to cause antithrombin-III deficiency which constitutes a strong risk factor for thrombosis. A reduction in the serum level of this protein is associated with severe cases of Coronavirus Disease 19 (COVID-19). [provided by RefSeq, Sep 2020]