

Product datasheet for **RC222893L3V**

ATP11C (NM_173694) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	ATP11C (NM_173694) Human Tagged ORF Clone Lentiviral Particle
Symbol:	ATP11C
Synonyms:	ATPIG; ATPIQ; HACXL
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_173694
ORF Size:	3396 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC222893).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_173694.3 , NP_775965.2
RefSeq Size:	6135 bp
RefSeq ORF:	3399 bp
Locus ID:	286410
UniProt ID:	Q8NB49
Cytogenetics:	Xq27.1
Protein Families:	Transmembrane
MW:	129.4 kDa



[View online »](#)

Gene Summary:

Catalytic component of a P4-ATPase flippase complex which catalyzes the hydrolysis of ATP coupled to the transport of aminophospholipids from the outer to the inner leaflet of various membranes and ensures the maintenance of asymmetric distribution of phospholipids. In the cell membrane of erythrocytes, it is required to maintain phosphatidylserine (PS) in the inner leaflet preventing its exposure on the surface. This asymmetric distribution is critical for the survival of erythrocytes in circulation since externalized PS is a phagocytic signal for splenic macrophages (PubMed:26944472). Phospholipid translocation seems also to be implicated in vesicle formation and in uptake of lipid signaling molecules (By similarity). Required for B cell differentiation past the pro-B cell stage (By similarity). Seems to mediate PS flipping in pro-B cells (By similarity). May be involved in the transport of cholestatic bile acids (By similarity).[UniProtKB/Swiss-Prot Function]