

Product datasheet for RC222898L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRIK1 (NM_000830) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: GRIK1 (NM 000830) Human Tagged ORF Clone Lentiviral Particle

Symbol: GRIK1

Synonyms: EAA3; EEA3; GLR5; GluK1; gluR-5; GLUR5

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 000830

ORF Size: 2754 bp

ORF Nucleotide

TI ODE

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC222898).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000830.3

 RefSeq Size:
 3231 bp

 RefSeq ORF:
 2757 bp

 Locus ID:
 2897

 UniProt ID:
 P39086

Cytogenetics: 21q21.3

Domains: lig_chan, ANF_receptor

Protein Families: Druggable Genome, Ion Channels: Glutamate Receptors, Transmembrane





GRIK1 (NM_000830) Human Tagged ORF Clone Lentiviral Particle - RC222898L1V

Protein Pathways: Neuroactive ligand-receptor interaction

MW: 104 kDa

Gene Summary: Glutamate receptors are the predominant excitatory neurotransmitter receptors in the

mammalian brain and are activated in a variety of normal neurophysiologic processes. This gene product belongs to the kainate family of glutamate receptors, which are composed of four subunits and function as ligand-activated ion channels. The subunit encoded by this gene is subject to RNA editing (CAG->CGG; Q->R) within the second transmembrane domain, which is thought to alter the properties of ion flow. Alternative splicing, resulting in transcript variants encoding different isoforms, has been noted for this gene. [provided by RefSeq, Jul

2008]