

Product datasheet for RC222952L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DNA Ligase III (LIG3) (NM 002311) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: DNA Ligase III (LIG3) (NM 002311) Human Tagged ORF Clone Lentiviral Particle

Symbol: LIG3

Synonyms: LIG2; LIG3alpha

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_002311 **ORF Size:** 2847 bp

ORF Nucleotide

- -

Sequence:
OTI Disclaimer:

Cytogenetics:

The ORF insert of this clone is exactly the same as(RC222952).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002311.4, NP 002302.2

17q12

 RefSeq Size:
 3103 bp

 RefSeq ORF:
 2850 bp

 Locus ID:
 3980

 UniProt ID:
 P49916

Domains: DNA_ligase, zf-PARP

Protein Families: Druggable Genome





Protein Pathways: Base excision repair

MW: 106 kDa

Gene Summary: This gene is a member of the DNA ligase family. Each member of this family encodes a

protein that catalyzes the joining of DNA ends but they each have a distinct role in DNA metabolism. The protein encoded by this gene is involved in excision repair and is located in both the mitochondria and nucleus, with translation initiation from the upstream start codon allowing for transport to the mitochondria and translation initiation from a downstream start codon allowing for transport to the nucleus. Additionally, alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]