

## Product datasheet for **RC222992**

### G protein alpha S (GNAS) (NM\_001077488) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	G protein alpha S (GNAS) (NM_001077488) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	G protein alpha S
Synonyms:	AHO; C20orf45; GNAS1; GPSA; GSA; GSP; NESP; PITA3; POH; SCG6; SgVI
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC222992 representing NM_001077488 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGGCTGCCTCGGGAACAGTAAGACCGAGGACCAGCGCAACGAGGAGAAGGCGCAGCGTGAGGCCAACA  
AAAAGATCGAGAAGCAGCTGCAGAAGGACAAGCAGGTCTACGGGCCACGCACCGCTGCTGCTGCTGGG  
TGCTGGAGAATCTGGTAAAAGCACCATTGTGAAGCAGATGAGGATCCTGCATGTTAATGGGTTAATGGA  
GAGGGCGGCGAAGAGGACCCGACAGGCTGCAAGGAGCAACAGCGATGGCAGTGAGAAGGCAACCAAAGTGC  
AGGACATCAAAAACAACCTGAAAGAGGCGATTGAAACCATTGTGGCCGCATGAGCAACCTGGTGCCCCC  
CGTGGAGCTGGCCAAACCCGAGAACCAGTTCAGAGTGGACTACATCCTGAGTGTGATGAACGTGCCTGAC  
TTTGACTTCCCTCCCGAATTCTATGAGCATGCCAAGGCTCTGTGGGAGGATGAAGGAGTGCCTGCCTGCT  
ACGAACGCTCCAACGAGTACCAGCTGATTGACTGTGCCAGTACTTCTGGACAAGATCGACGTGATCAA  
GCAGGCTGACTATGTGCCGAGCGATCAGGACCTGCTTCGCTGCCGTGTCCTGACTTCTGGAATCTTTGAG  
ACCAAGTTCAGGTGGACAAAGTCAACTCCACATGTTTGACGTGGGTGGCCAGCGCGATGAACGCCGCA  
AGTGGATCCAGTGCTTCAACGATGTGACTGCCATCATCTCGTGGTGGCCAGCAGCAGCTACAACATGGT  
CATCCGGGAGGACAACCAGCAACCGCCTGCAGGAGGCTCTGAACCTCTTCAAGAGCATCTGGAACAAC  
AGATGGCTGCGCACCATCTGTGATCCTGTTCCCTCAACAAGCAAGATCTGCTCGTGAGAAAGTCTTTG  
CTGGGAAATCGAAGATTGAGGACTACTTCCAGAATTTGCTCGCTACACTACTCCTGAGGATGCTACTCC  
CGAGCCCGGAGAGGACCCACGCGTGACCCGGGCAAGTACTTCATTTCGAGATGAGTTTCTGAGGATCAGC  
ACTGCCAGTGGAGATGGGCGTCACTACTGCTACCTCATTTACCTGCGTGTGGACACTGAGAATATCC  
GCCGTGTGTTCAACGACTGCCGTGACATCATTACGCGCATGCACCTTCGTCAGTACGAGCTGCTC

AG**CGGACCG**ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC  
TGGATTACAAGGATGACGACGATAAGGTTTAA



[View online >](#)

**Protein Sequence:** >RC222992 representing NM\_001077488  
 Red=Cloning site Green=Tags(s)


MGCLGNSKTEDQRNEEKAQREANKKIEKQLQKDKQVYRATHRLLLLGAGESGKSTIVKQMRILHVNGFNG  
 EGGEEDPQAARSNSDGSEKATKVQDIKNNLKEAIETIVAAMSNL VPPVELANPENQFRVDYILSVMNVPD  
 FDFPPEFYEHAKALWEDEGVRACYERSNEYQLIDCAQYFLDKIDVIKQADYVPSDQDLLRRCVLTSGIFE  
 TKFQVDKVNFMFDVGGQRDERRKWIQCFNDVTAIIFVVAASSYNMVIREDNQTNRLQEALNLFKSIWNN  
 RWLRTISVILFLNKQDLLAEKVLGKSKIETYFPEFARYTTPEDATPEPGEDPRVTRAKYFIRDEFRLRIS  
 TASGDGRHYCYPHFTCAVDTENIRRVFNDCRDIIQRMHLRQYELL

SGP TRTRRLEQKLI SEEDLAANDILDYKDDDDKV

**Chromatograms:** [https://cdn.origene.com/chromatograms/mk8057\\_a06.zip](https://cdn.origene.com/chromatograms/mk8057_a06.zip)

**Restriction Sites:** SgfI-RsrII

**Cloning Scheme:**



**ACCN:** NM\_001077488


**ORF Size:** 1185 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at [custsupport@origene.com](mailto:custsupport@origene.com) or by calling 301.340.3188 option 3 for pricing and delivery.


The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001077488.5</a>
<b>RefSeq Size:</b>	1586 bp
<b>RefSeq ORF:</b>	1188 bp
<b>Locus ID:</b>	2778
<b>UniProt ID:</b>	<a href="#">O95467</a>
<b>Cytogenetics:</b>	20q13.32
<b>Protein Families:</b>	Druggable Genome, Secreted Protein
<b>Protein Pathways:</b>	Calcium signaling pathway, Dilated cardiomyopathy, Gap junction, GnRH signaling pathway, Long-term depression, Melanogenesis, Taste transduction, Vascular smooth muscle contraction, Vibrio cholerae infection
<b>MW:</b>	45.6 kDa
<b>Gene Summary:</b>	<p>This locus has a highly complex imprinted expression pattern. It gives rise to maternally, paternally, and biallelically expressed transcripts that are derived from four alternative promoters and 5' exons. Some transcripts contain a differentially methylated region (DMR) at their 5' exons, and this DMR is commonly found in imprinted genes and correlates with transcript expression. An antisense transcript is produced from an overlapping locus on the opposite strand. One of the transcripts produced from this locus, and the antisense transcript, are paternally expressed noncoding RNAs, and may regulate imprinting in this region. In addition, one of the transcripts contains a second overlapping ORF, which encodes a structurally unrelated protein - Alex. Alternative splicing of downstream exons is also observed, which results in different forms of the stimulatory G-protein alpha subunit, a key element of the classical signal transduction pathway linking receptor-ligand interactions with the activation of adenylyl cyclase and a variety of cellular responses. Multiple transcript variants encoding different isoforms have been found for this gene. Mutations in this gene result in pseudohypoparathyroidism type 1a, pseudohypoparathyroidism type 1b, Albright hereditary osteodystrophy, pseudopseudohypoparathyroidism, McCune-Albright syndrome, progressive osseus heteroplasia, polyostotic fibrous dysplasia of bone, and some pituitary tumors. [provided by RefSeq, Aug 2012]</p>

Product images:



Circular map for RC222992



Western blot validation of overexpression lysate (Cat# [LY421439]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC222992 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).