

Product datasheet for **RC222995L3V**

Cytochrome P450 2A6 (CYP2A6) (NM_000762) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Cytochrome P450 2A6 (CYP2A6) (NM_000762) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Cytochrome P450 2A6
Synonyms:	CPA6; CYP2A; CYP2A3; CYP11A6; P450C2A; P450PB
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_000762
ORF Size:	1482 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC222995).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_000762.4
RefSeq Size:	1775 bp
RefSeq ORF:	1485 bp
Locus ID:	1548
UniProt ID:	P11509
Cytogenetics:	19q13.2
Domains:	p450
Protein Families:	Druggable Genome, P450, Transmembrane


[View online »](#)

Protein Pathways:	Caffeine metabolism, Drug metabolism - cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Retinol metabolism
MW:	56.5 kDa
Gene Summary:	This gene, CYP2A6, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to hydroxylate coumarin, and also metabolizes nicotine, aflatoxin B1, nitrosamines, and some pharmaceuticals. Individuals with certain allelic variants are said to have a poor metabolizer phenotype, meaning they do not efficiently metabolize coumarin or nicotine. This gene is part of a large cluster of cytochrome P450 genes from the CYP2A, CYP2B and CYP2F subfamilies on chromosome 19q. The gene was formerly referred to as CYP2A3; however, it has been renamed CYP2A6. [provided by RefSeq, Jul 2008]