

## Product datasheet for RC223000L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NFAT4 (NFATC3) (NM\_173163) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: NFAT4 (NFATC3) (NM 173163) Human Tagged ORF Clone Lentiviral Particle

Symbol: NFATC3

Synonyms: NF-AT4c; NFAT4; NFATX

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_173163 **ORF Size:** 3195 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC223000).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 173163.2

 RefSeq Size:
 6438 bp

 RefSeq ORF:
 3198 bp

 Locus ID:
 4775

 UniProt ID:
 Q12968

 Cytogenetics:
 16q22.1

**Protein Families:** Druggable Genome, Transcription Factors





## NFAT4 (NFATC3) (NM\_173163) Human Tagged ORF Clone Lentiviral Particle - RC223000L4V

**Protein Pathways:** Axon guidance, B cell receptor signaling pathway, Natural killer cell mediated cytotoxicity, T

cell receptor signaling pathway, VEGF signaling pathway, Wnt signaling pathway

MW: 115.1 kDa

**Gene Summary:** The product of this gene is a member of the nuclear factors of activated T cells DNA-binding

transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation and an inducible nuclear component. Other members of this family participate to form this complex also. The product of this gene plays a role in the regulation of gene expression in T cells and immature thymocytes. Several transcript variants encoding distinct isoforms have

been identified for this gene. [provided by RefSeq, Nov 2010]