

Product datasheet for RC223097L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CYP2F1 (NM_000774) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CYP2F1 (NM 000774) Human Tagged ORF Clone Lentiviral Particle

Symbol: CYP2F1

Synonyms: C2F1; CYP2F; CYPIIF1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_000774

ORF Size: 1473 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC223097).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000774.3</u>

 RefSeq Size:
 1826 bp

 RefSeq ORF:
 1476 bp

 Locus ID:
 1572

 UniProt ID:
 P24903

 Cytogenetics:
 19q13.2

Protein Families: Druggable Genome, P450, Transmembrane

Protein Pathways: Metabolism of xenobiotics by cytochrome P450





ORIGENE

MW: 55.3 kDa

Gene Summary: This gene encodes a member of the cytochrome P450 superfamily of enzymes. The

cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes

to the endoplasmic reticulum and is known to dehydrogenate 3-methylindole, an endogenous toxin derived from the fermentation of tryptophan, as well as xenobiotic substrates such as naphthalene and ethoxycoumarin. This gene is part of a large cluster of cytochrome P450 genes from the CYP2A, CYP2B and CYP2F subfamilies on chromosome 19q.

[provided by RefSeq, Jul 2008]