

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC223226L3V

OAS2 (NM_001032731) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	OAS2 (NM_001032731) Human Tagged ORF Clone Lentiviral Particle
Symbol:	OAS2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001032731
ORF Size:	516 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC223226).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001032731.1</u>
RefSeq Size:	2123 bp
RefSeq ORF:	519 bp
Locus ID:	4939
UniProt ID:	<u>P29728</u>
Cytogenetics:	12q24.13
Protein Families:	Druggable Genome
MW:	20 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes a member of the 2-5A synthetase family, essential proteins involved in the
innate immune response to viral infection. The encoded protein is induced by interferons
and uses adenosine triphosphate in 2'-specific nucleotidyl transfer reactions to synthesize
2',5'-oligoadenylates (2-5As). These molecules activate latent RNase L, which results in viral
RNA degradation and the inhibition of viral replication. The three known members of this
gene family are located in a cluster on chromosome 12. Alternatively spliced transcript
variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US