

Product datasheet for RC223260L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PLAGL1 (NM_006718) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PLAGL1 (NM_006718) Human Tagged ORF Clone Lentiviral Particle

Symbol: PLAGL1

Synonyms: LOT1; ZAC; ZAC1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_006718 **ORF Size:** 1389 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC223260).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 006718.3

RefSeq Size: 3411 bp
RefSeq ORF: 1392 bp
Locus ID: 5325
UniProt ID: Q9UM63

Cytogenetics: 6q24.2 Domains: zf-C2H2

Protein Families: Transcription Factors



PLAGL1 (NM_006718) Human Tagged ORF Clone Lentiviral Particle - RC223260L4V

MW: 50.8 kDa

Gene Summary: This gene encodes a C2H2 zinc finger protein that functions as a suppressor of cell growth.

This gene is often deleted or methylated and silenced in cancer cells. In addition,

overexpression of this gene during fetal development is thought to be the causal factor for transient neonatal diabetes mellitus (TNDM). Alternative splicing and the use of alternative promoters results in multiple transcript variants encoding two different protein isoforms. The P1 downstream promoter of this gene is imprinted, with preferential expression from the

paternal allele in many tissues. [provided by RefSeq, Nov 2015]