

#### OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC223317L1

### Angiotensin II Type 2 Receptor (AGTR2) (NM\_000686) Human Tagged Lenti ORF Clone

#### **Product data:**

| Product Type:                | Expression Plasmids   |
|------------------------------|---|
| Product Name:                | Angiotensin II Type 2 Receptor (AGTR2) (NM_000686) Human Tagged Lenti ORF Clone |
| Tag:                         | Myc-DDK   |
| Symbol:                      | Angiotensin II Type 2 Receptor  |
| Synonyms:                    | AT2; ATGR2; MRX88   |
| Mammalian Cell<br>Selection: | None  |
| Vector:                      | pLenti-C-Myc-DDK (PS100064)   |
| E. coli Selection:           | Chloramphenicol (34 ug/mL)  |
| ORF Nucleotide<br>Sequence:  | The ORF insert of this clone is exactly the same as(RC223317).                  |
| <b>Restriction Sites:</b>    | Sgfl-Mlul   |
| Cloning Scheme:              |   |
|                              | Cloning sites used for ORF Shuttling:   |
|                              | Sgf1         ORF         Mlu I            GCG ATC GC         ATG // NNÑ ACG CGT |



ACCN: ORF Size:

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

NM\_000686


1089 bp

| Angiotensin II Type 2 Receptor (AGTR2) (NM_000686) Human Tagged Lenti ORF Clone –<br>RC223317L1 |   |
|---|---|
| OTI Disclaimer:   | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>   |
| OTI Annotation:   | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.  |
| Components:   | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).  |
| Reconstitution Me   | <ul> <li>2. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ul>   |
| RefSeq:   | <u>NM 000686.3</u>  |
| RefSeq Size:  | 2448 bp   |
| RefSeq ORF:   | 1092 bp   |
| Locus ID:   | 186   |
| UniProt ID:   | <u>P50052</u>   |
| Cytogenetics:   | Xq23  |
| Protein Families:   | Druggable Genome, GPCR, Transmembrane   |
| Protein Pathways:   | Neuroactive ligand-receptor interaction, Renin-angiotensin system   |
| MW:   | 41 kDa  |
| Gene Summary:   | The protein encoded by this gene belongs to the G-protein coupled receptor 1 family, and<br>functions as a receptor for angiotensin II. It is an intergral membrane protein that is highly<br>expressed in fetus and in neonates, but scantily in adult tissues, except brain, adrenal<br>medulla, and atretic ovary. This receptor has been shown to mediate programmed cell death<br>and this apoptotic function may play an important role in developmental biology and<br>pathophysiology. Mutations in this gene are been associated with X-linked cognitive disability.<br>Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV) and SARS-CoV-2 infection results<br>in down-regulation of angiotensin converting enzyme-2 (ACE2) receptors, the effects of which,<br>triggers serious inflammatory lesions in the tissues involved, primarily in the lungs. The<br>inflammatory reaction appears to be mediated by angiotensin II derivatives, including the<br>angiotensin AT2 receptor which has been found to be upregulated in bronchoalveolar lavage<br>samples from Coronavirus disease 2019 (COVID19) patients. [provided by RefSeq, Jul 2020] |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



## **Product images:**



Circular map for RC223317L1

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US