

Product datasheet for RC223373L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PARP2 (NM_005484) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PARP2 (NM 005484) Human Tagged ORF Clone Lentiviral Particle

Symbol: PARP2

Synonyms: ADPRTL2; ADPRTL3; ARTD2; pADPRT-2; PARP-2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_005484

 ORF Size:
 1749 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC223373).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005484.3

RefSeq Size: 1904 bp
RefSeq ORF: 1752 bp
Locus ID: 10038
UniProt ID: Q9UGN5
Cytogenetics: 14q11.2

Domains: PARP, PARP_reg

Protein Families: Druggable Genome





PARP2 (NM_005484) Human Tagged ORF Clone Lentiviral Particle - RC223373L3V

Protein Pathways: Base excision repair

MW: 66 kDa

Gene Summary: This gene encodes poly(ADP-ribosyl)transferase-like 2 protein, which contains a catalytic

domain and is capable of catalyzing a poly(ADP-ribosyl)ation reaction. This protein has a catalytic domain which is homologous to that of poly (ADP-ribosyl) transferase, but lacks an N-terminal DNA binding domain which activates the C-terminal catalytic domain of poly (ADP-ribosyl) transferase. The basic residues within the N-terminal region of this protein may bear potential DNA-binding properties, and may be involved in the nuclear and/or nucleolar targeting of the protein. Two alternatively spliced transcript variants encoding distinct

isoforms have been found. [provided by RefSeq, Jul 2008]