

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC223407L4V

## USP37 (NM\_020935) Human Tagged ORF Clone Lentiviral Particle

## Product data:

Product Type:	Lentiviral Particles
Product Name:	USP37 (NM_020935) Human Tagged ORF Clone Lentiviral Particle
Symbol:	USP37
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_020935
ORF Size:	2937 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC223407).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 020935.1</u>
RefSeq Size:	8015 bp
RefSeq ORF:	2940 bp
Locus ID:	57695
UniProt ID:	<u>Q86T82</u>
Cytogenetics:	2q35
Protein Families:	Protease
MW:	110 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Deubiquitinase that antagonizes the anaphase-promoting complex (APC/C) during G1/S<br/>transition by mediating deubiquitination of cyclin-A (CCNA1 and CCNA2), thereby promoting S<br/>phase entry. Specifically mediates deubiquitination of 'Lys-11'-linked polyubiquitin chains, a<br/>specific ubiquitin-linkage type mediated by the APC/C complex. Also mediates<br/>deubiquitination of 'Lys-48'-linked polyubiquitin chains in vitro. Phosphorylation at Ser-628<br/>during G1/S phase maximizes the deubiquitinase activity, leading to prevent degradation of<br/>cyclin-A (CCNA1 and CCNA2) (PubMed:21596315). Plays an important role in the regulation of<br/>DNA replication by stabilizing the licensing factor CDT1 (PubMed:27296872).<br/>[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US