

Product datasheet for RC223438L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RNF14 (NM_183398) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RNF14 (NM_183398) Human Tagged ORF Clone Lentiviral Particle

Symbol: RNF14

Synonyms: ARA54; HFB30; HRIHFB2038; TRIAD2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 183398

ORF Size: 1425 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC223438).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 183398.2, NP 899645.1

RefSeq Size: 4243 bp
RefSeq ORF: 1047 bp
Locus ID: 9604
UniProt ID: Q9UBS8

Cytogenetics: 5q31.3

Protein Families: Druggable Genome, Transcription Factors

MW: 53.8 kDa







Gene Summary:

The protein encoded by this gene contains a RING zinc finger, a motif known to be involved in protein-protein interactions. This protein interacts with androgen receptor (AR) and may function as a coactivator that induces AR target gene expression in prostate. A dominant negative mutant of this gene has been demonstrated to inhibit the AR-mediated growth of prostate cancer. This protein also interacts with class III ubiquitin-conjugating enzymes (E2s) and may act as a ubiquitin-ligase (E3) in the ubiquitination of certain nuclear proteins. Six alternatively spliced transcript variants encoding two distinct isoforms have been reported. [provided by RefSeq, Jan 2011]