

Product datasheet for RC223519L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Melanopsin (OPN4) (NM_033282) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Melanopsin (OPN4) (NM_033282) Human Tagged ORF Clone Lentiviral Particle

Symbol: Melanopsin

Synonyms: MOP

Mammalian Cell None

Selection: Vector:

pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_033282

ORF Size: 1434 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC223519).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 033282.2</u>

RefSeq Size: 2386 bp
RefSeq ORF: 1437 bp
Locus ID: 94233
UniProt ID: Q9UHM6
Cytogenetics: 10q23.2

Protein Families: Druggable Genome, Transmembrane

MW: 52.7 kDa





Gene Summary:

Opsins are members of the guanine nucleotide-binding protein (G protein)-coupled receptor superfamily. This gene encodes a photoreceptive opsin protein that is expressed within the ganglion and amacrine cell layers of the retina. In mouse, retinal ganglion cell axons expressing this gene projected to the suprachiasmatic nucleus and other brain nuclei involved in circadian photoentrainment. In mouse, this protein is coupled to a transient receptor potential (TRP) ion channel through a G protein signaling pathway and produces a physiologic light response via membrane depolarization and increased intracellular calcium. The protein functions as a sensory photopigment and may also have photoisomerase activity. Experiments with knockout mice indicate that this gene attenuates, but does not abolish, photoentrainment. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]