

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC223543L3V

MAFK (NM_002360) Human Tagged ORF Clone Lentiviral Particle

Product data:

Lentiviral Particles
MAFK (NM_002360) Human Tagged ORF Clone Lentiviral Particle
MAFK
NFE2U; P18
Puromycin
pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Myc-DDK
NM_002360
468 bp
The ORF insert of this clone is exactly the same as(RC223543).
The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<u>NM 002360.2</u>
1581 bp
471 bp
7975
<u>060675</u>
7p22.3
bZIP_Maf, BRLZ
Druggable Genome, Transcription Factors



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	MAFK (NM_002360) Human Tagged ORF Clone Lentiviral Particle – RC223543L3V
MW:	17.3 kDa
Gene Summary:	The developmentally regulated expression of the globin genes depends on upstream regulatory elements termed locus control regions (LCRs). LCRs are associated with powerful enhancer activity that is mediated by the transcription factor NFE2 (nuclear factor erythroid-2). NFE2 recognition sites are also present in the gene promoters of 2 heme biosynthetic enzymes, porphobilinogen deaminase (PBGD; MIM 609806) and ferrochelatase (FECH; MIM 612386). NFE2 DNA-binding activity consists of a heterodimer containing an 18-kD Maf protein (MafF, MafG (MIM 602020), or MafK) and p45 (MIM 601490). Both subunits are members of the activator protein-1 superfamily of basic leucine zipper (bZIP) proteins (see MIM 165160). Maf homodimers suppress transcription at NFE2 sites.[supplied by OMIM, Nov 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US