

## Product datasheet for RC223545L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **UGT2A3 (NM\_024743) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** UGT2A3 (NM\_024743) Human Tagged ORF Clone Lentiviral Particle

Symbol: UGT2A3

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_024743

ORF Size: 1581 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC223545).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 024743.2, NP 079019.2</u>

 RefSeq Size:
 2974 bp

 RefSeq ORF:
 1584 bp

 Locus ID:
 79799

UniProt ID: Q6UWM9

Cytogenetics: 4q13.2

**Protein Families:** Transmembrane



## UGT2A3 (NM\_024743) Human Tagged ORF Clone Lentiviral Particle - RC223545L3V

Protein Pathways: Androgen and estrogen metabolism, Ascorbate and aldarate metabolism, Drug metabolism -

cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Pentose and glucuronate interconversions, Porphyrin and

chlorophyll metabolism, Retinol metabolism, Starch and sucrose metabolism

**MW:** 60.1 kDa

Gene Summary: UDP-glucuronosyltransferases catalyze phase II biotransformation reactions in which

lipophilic substrates are conjugated with glucuronic acid to increase water solubility and enhance excretion. They are of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds (By similarity).

[UniProtKB/Swiss-Prot Function]