

## Product datasheet for RC223572L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **GAK (NM\_005255) Human Tagged ORF Clone Lentiviral Particle**

## **Product data:**

Product Type: Lentiviral Particles

**Product Name:** GAK (NM\_005255) Human Tagged ORF Clone Lentiviral Particle

Symbol: GAK

Synonyms: DNAJ26; DNAJC26

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 005255

ORF Size: 3933 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC223572).

\_\_\_\_

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 005255.1

 RefSeq Size:
 4331 bp

 RefSeq ORF:
 3936 bp

 Locus ID:
 2580

 UniProt ID:
 014976

 Cytogenetics:
 4p16.3

**Domains:** pkinase, TyrKc, DnaJ, S\_TKc

**Protein Families:** Druggable Genome, Protein Kinase



ORIGENE

MW: 143 kDa

**Gene Summary:** 

In all eukaryotes, the cell cycle is governed by cyclin-dependent protein kinases (CDKs), whose activities are regulated by cyclins and CDK inhibitors in a diverse array of mechanisms that involve the control of phosphorylation and dephosphorylation of Ser, Thr or Tyr residues. Cyclins are molecules that possess a consensus domain called the 'cyclin box.' In mammalian cells, 9 cyclin species have been identified, and they are referred to as cyclins A through I. Cyclin G is a direct transcriptional target of the p53 tumor suppressor gene product and thus functions downstream of p53. GAK is an association partner of cyclin G and CDK5. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2015]