

## Product datasheet for RC223635L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ATP7B (NM\_000053) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** ATP7B (NM\_000053) Human Tagged ORF Clone Lentiviral Particle

Symbol: ATP7B

Synonyms: PWD; WC1; WD; WND

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_000053 **ORF Size:** 4395 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC223635).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000053.2

 RefSeq Size:
 6644 bp

 RefSeq ORF:
 4398 bp

 Locus ID:
 540

 UniProt ID:
 P35670

 Cytogenetics:
 13q14.3

**Domains:** E1-E2\_ATPase, Hydrolase, HMA

**Protein Families:** Druggable Genome, Transmembrane





ORIGENE

**MW:** 157.1 kDa

**Gene Summary:** 

This gene is a member of the P-type cation transport ATPase family and encodes a protein with several membrane-spanning domains, an ATPase consensus sequence, a hinge domain, a phosphorylation site, and at least 2 putative copper-binding sites. This protein is a monomer, and functions as a copper-transporting ATPase which exports copper out of the cells, such as the efflux of hepatic copper into the bile. Alternate transcriptional splice variants, encoding different isoforms with distinct cellular localizations, have been characterized. Mutations in this gene have been associated with Wilson disease which is characterized by copper accumulation. [provided by RefSeq, Dec 2019]