

Product datasheet for **RC223737L4V**

Bcl 7A (BCL7A) (NM_020993) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Bcl 7A (BCL7A) (NM_020993) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Bcl 7A
Synonyms:	BCL7
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_020993
ORF Size:	693 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC223737).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_020993.3 , NP_066273.1
RefSeq Size:	3714 bp
RefSeq ORF:	696 bp
Locus ID:	605
UniProt ID:	Q4VC05
Cytogenetics:	12q24.31
Domains:	BCL_N
Protein Families:	Druggable Genome



[View online »](#)

MW: 24.8 kDa

Gene Summary: This gene is directly involved, with Myc and IgH, in a three-way gene translocation in a Burkitt lymphoma cell line. As a result of the gene translocation, the N-terminal region of the gene product is disrupted, which is thought to be related to the pathogenesis of a subset of high-grade B cell non-Hodgkin lymphoma. The N-terminal segment involved in the translocation includes the region that shares a strong sequence similarity with those of BCL7B and BCL7C. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]