

# Product datasheet for RC223780L4


# FADS2 (NM\_004265) Human Tagged Lenti ORF Clone

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	FADS2 (NM_004265) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	FADS2
Synonyms:	D6D; DES6; FADSD6; LLCDL2; SLL0262; TU13
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC223780).
<b>Restriction Sites:</b>	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf1         ORF         Mlu1            GCG ATC GC         ATG          NNN         ACG CGT



ACCN: ORF Size: NM\_004265 1332 bp



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


<b>Gene</b> FADS2 (NM_004265) Human Tagged Lenti ORF Clone – RC223780L4	
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Meth	<ul> <li>od: 1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ul>
RefSeq:	<u>NM 004265.2</u>
RefSeq Size:	3149 bp
RefSeq ORF:	1335 bp
Locus ID:	9415
UniProt ID:	<u>O95864</u>
Cytogenetics:	11q12.2
Domains:	heme_1, FA_desaturase
Protein Families:	Transmembrane
Protein Pathways:	alpha-Linolenic acid metabolism, Biosynthesis of unsaturated fatty acids, PPAR signaling pathway
MW:	52.1 kDa

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### SADS2 (NM\_004265) Human Tagged Lenti ORF Clone – RC223780L4

Gene Summary:The protein encoded by this gene is a member of the fatty acid desaturase (FADS) gene<br/>family. Desaturase enzymes regulate unsaturation of fatty acids through the introduction of<br/>double bonds between defined carbons of the fatty acyl chain. FADS family members are<br/>considered fusion products composed of an N-terminal cytochrome b5-like domain and a C-<br/>terminal multiple membrane-spanning desaturase portion, both of which are characterized<br/>by conserved histidine motifs. This gene is clustered with family members at 11q12-q13.1; this<br/>cluster is thought to have arisen evolutionarily from gene duplication based on its similar<br/>exon/intron organization. Alternative splicing results in multiple transcript variants encoding<br/>different isoforms. [provided by RefSeq, Jul 2013]

### **Product images:**



Circular map for RC223780L4

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US