

Product datasheet for RC223917L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Deoxyguanosine kinase (DGUOK) (NM_080918) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Deoxyguanosine kinase (DGUOK) (NM_080918) Human Tagged ORF Clone Lentiviral Particle

Symbol: Deoxyguanosine kinase

Synonyms: dGK; MTDPS3; NCPH; PEOB4

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 080918

ORF Size: 567 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC223917).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 080918.1

 RefSeq Size:
 880 bp

 RefSeq ORF:
 570 bp

 Locus ID:
 1716

 UniProt ID:
 Q16854

 Cytogenetics:
 2p13.1

Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Purine metabolism





Deoxyguanosine kinase (DGUOK) (NM_080918) Human Tagged ORF Clone Lentiviral Particle – RC223917L3V

MW: 17.3 kDa

Gene Summary:

In mammalian cells, the phosphorylation of purine deoxyribonucleosides is mediated predominantly by two deoxyribonucleoside kinases, cytosolic deoxycytidine kinase and mitochondrial deoxyguanosine kinase. The protein encoded by this gene is responsible for phosphorylation of purine deoxyribonucleosides in the mitochondrial matrix. In addition, this protein phosphorylates several purine deoxyribonucleoside analogs used in the treatment of lymphoproliferative disorders, and this phosphorylation is critical for the effectiveness of the analogs. Alternative splice variants encoding different protein isoforms have been described for this gene. [provided by RefSeq, Jul 2008]