

# Product datasheet for RC223961

## EMA (MUC1) (NM\_001044393) Human Tagged ORF Clone

### **Product data:**

Product Type:	Expression Plasmids
Product Name:	EMA (MUC1) (NM_001044393) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	MUC1
Synonyms:	ADMCKD; ADMCKD1; ADTKD2; CA 15-3; CD227; EMA; H23AG; KL-6; MAM6; MCD; MCKD; MCKD1; MUC-1; MUC-1/SEC; MUC-1/X; MUC1/ZD; PEM; PEMT; PUM
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
ORF Nucleotide Sequence:	<pre>&gt;RC223961 representing NM_001044393 Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGACACCGGGCACCCAGTCTCCTTTCTTCCTGCTGCTGCTCCTCACAGTGCTTACAGTTGTTACGGGTT CTGGTCATGCAAGCTCTACCCCAGGTGGAGAAAAGGAGACTTCGGCTACCCAGAGAAGTTCAGTGCCCAG CTCTACTGAGAAGAATGCTATCCCAGCACCGACTACTACCAAGAGCTGCAGAGAGAG
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG <b>GTTTAA</b>
Protein Sequence:	<pre>&gt;RC223961 representing NM_001044393 Red=Cloning site Green=Tags(s)</pre>
	MTPGTQSPFFLLLLLTVLTVVTGSGHASSTPGGEKETSATQRSSVPSSTEKNAIPAPTTTKSCRETFLKC FCRFINKGVFWASPILSSVWGWGARLGHRAAGAGLCSGCAGHCLSHCLGCLSVPPKELRAAGHLSSPGYL PSYERVPHLPHPWALCAP
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Restriction Sites:	Sgfl-Mlul



View online »


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn




#### **Cloning Scheme:**



\* The last codon before the Stop codon of the ORF

#### Plasmid Map:



ACCN:	NM_001044393
ORF Size:	474 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

SRIGENE EMA (MUC1) (NM_001044393) Human Tagged ORF Clone – RC223961	
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Metho	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 001044393.3</u>
RefSeq Size:	978 bp
RefSeq ORF:	477 bp
Locus ID:	4582
UniProt ID:	<u>P15941</u>
Cytogenetics:	1q22
Protein Families:	Druggable Genome, Secreted Protein, Transmembrane
MW:	16.6 kDa
Gene Summary:	This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly

multiple transcript variants.[provided by RefSeq, Feb 2011]

polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US