

## Product datasheet for RC223966L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ERCC6L (NM\_017669) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: ERCC6L (NM\_017669) Human Tagged ORF Clone Lentiviral Particle

Symbol: ERCC6L

Synonyms: PICH; RAD26L

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_017669 **ORF Size:** 3750 bp

**ORF Nucleotide** 

'

Sequence:

The ORF insert of this clone is exactly the same as(RC223966).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 017669.2</u>

 RefSeq Size:
 4224 bp

 RefSeq ORF:
 3753 bp

 Locus ID:
 54821

 UniProt ID:
 Q2NKX8

 Cytogenetics:
 Xq13.1

MW: 141.1 kDa







## **Gene Summary:**

This gene encodes a member of the SWItch/Sucrose Non-Fermentable (SWI/SNF2) family of proteins, and contains a SNF2-like ATPase domain and a PICH family domain. One distinguishing feature of this SWI/SNF protein family member is that during interphase, the protein is excluded from the nucleus, and only associates with chromatin after the nuclear envelope has broken down. This protein is a DNA translocase that is thought to bind double-stranded DNA that is exposed to stretching forces, such as those exerted by the mitotic spindle. This protein associates with ribosomal DNA and ultra-fine DNA bridges (UFBs), fine structures that connect sister chromatids during anaphase at some sites such as fragile sites, telomeres and centromeres. This gene is required for the faithful segregation of sister chromatids during mitosis, and the ATPase activity of this protein required for the resolution of UFBs before cytokinesis. [provided by RefSeq, May 2017]