

Product datasheet for RC223979L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HLA (HLA-DQA2) (NM 020056) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: HLA (HLA-DQA2) (NM 020056) Human Tagged ORF Clone Lentiviral Particle

Symbol: HLA-DQA2

Synonyms: DC-alpha; DX-ALPHA; HLA-DCA; HLA-DXA; HLADQA2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_020056

ORF Size: 765 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC223979).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 020056.2, NP 064440.1

 RefSeq Size:
 1709 bp

 RefSeq ORF:
 768 bp

 Locus ID:
 3118

 UniProt ID:
 P01906

 Cytogenetics:
 6p21.32

Domains: MHC_II_alpha, ig, IGc1

Protein Families: Transmembrane





HLA (HLA-DQA2) (NM_020056) Human Tagged ORF Clone Lentiviral Particle - RC223979L4V

Protein Pathways: Allograft rejection, Antigen processing and presentation, Asthma, Autoimmune thyroid

disease, Cell adhesion molecules (CAMs), Graft-versus-host disease, Systemic lupus

erythematosus, Type I diabetes mellitus, Viral myocarditis

MW: 27.9 kDa

Gene Summary: This gene belongs to the HLA class II alpha chain family. The encoded protein forms a

heterodimer with a class II beta chain. It is located in intracellular vesicles and plays a central role in the peptide loading of MHC class II molecules by helping to release the CLIP molecule from the peptide binding site. Class II molecules are expressed in antigen presenting cells (B lymphocytes, dendritic cells, macrophages) and are used to present antigenic peptides on the

cell surface to be recognized by CD4 T-cells. [provided by RefSeq, Jun 2010]