

## Product datasheet for RC224009L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## LEC (CCL16) (NM\_004590) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** LEC (CCL16) (NM\_004590) Human Tagged ORF Clone Lentiviral Particle

Symbol: LEC

Synonyms: CKb12; HCC-4; ILINCK; LCC-1; LEC; LMC; Mtn-1; NCC-4; NCC4; SCYA16; SCYL4

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 004590

ORF Size: 360 bp

**ORF Nucleotide** 

TI ODE '

Sequence:

The ORF insert of this clone is exactly the same as(RC224009).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 004590.2</u>

 RefSeq Size:
 1497 bp

 RefSeq ORF:
 363 bp

 Locus ID:
 6360

 UniProt ID:
 015467

 Cytogenetics:
 17q12

Domains: IL8

**Protein Families:** Druggable Genome, Secreted Protein





## LEC (CCL16) (NM\_004590) Human Tagged ORF Clone Lentiviral Particle - RC224009L1V

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction

**MW:** 13.6 kDa

**Gene Summary:** This gene is one of several cytokine genes clustered on the q-arm of chromosome 17.

Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for lymphocytes and monocytes but not for neutrophils. This cytokine also shows a potent myelosuppressive activity and suppresses proliferation of myeloid progenitor cells. The expression of this gene is

upregulated by IL-10. [provided by RefSeq, Jul 2008]